

Large stellar disks in small elliptical galaxies

Hans-Walter Rix^{1,4}, Marcella Carollo^{2,5} and Ken Freeman³

Astrophysical Journal Letters, March 1999.

¹ Steward Observatory

² Johns Hopkins University

³ Mount Stromlo and Siding Springs Observatory

⁴ Alfred P. Sloan Fellow

⁵ Hubble Fellow

ABSTRACT

We present the rotation velocities V and velocity dispersions σ along the principal axes of seven elliptical galaxies less luminous than $M_B = -19.5$. These kinematics extend beyond the half-light radii for all systems in this photometrically selected sample. At large radii the kinematics not only confirm that rotation and “diskiness” are important in faint ellipticals, as was previously known, but also demonstrate that in most sample galaxies the stars at large galactocentric distances have $(V/\sigma)_{max} \sim 2$, similar to the disks in *bona-fide* S0 galaxies. Comparing this high degree of ordered stellar motion in all sample galaxies with numerical simulations of dissipationless mergers argues against mergers with mass ratios $\leq 3 : 1$ as an important mechanism in the final shaping of low-luminosity ellipticals, and favors instead the dissipative formation of a disk.

subject headings: galaxies: formation - galaxies: evolution - galaxies: structure - galaxies: elliptical and lenticular - galaxies: kinematics and dynamics

1. Introduction

Elliptical galaxies occupy only a small volume in the stable parameter space defined by mass, luminosity, compactness, and rotational support (*e.g.* Kormendy and Djorgovski 1989; de Zeeuw and Franx 1991). This means that their present-day structure is not determined by stability, but by their formation history, specifically by the relative importance and the time ordering of dissipative processes *vs.* merging. Yet, the actual role of gas dissipation, resulting in gaseous and stellar disks, and of violent relaxation, leading to spheroidal systems with largely random stellar motions, is still under debate for ellipticals of different luminosity classes.

Observationally it has been established that “low-luminosity” ellipticals ($L \lesssim \frac{1}{2}L_*$) show more rotation and photometric diskiness than the brighter systems (*e.g.*, Davies *et al.* 1983; Bender *et al.* 1989). Yet, taken alone, these results cannot constrain uniquely their formation history, as there are many paths towards systems with modest rotation. For example, N-body simulations of dissipationless major mergers can produce rotating remnants $(V/\sigma)_{max} \lesssim 1$ with disk-like isophotes (*e.g.*, Heyl, Hernquist & Spergel 1996; Weil & Hernquist 1996). On the basis of statistics, Rix & White (1990) argued that most of the so-called faint ellipticals are not just “disky” objects, but are rather the face-on counterparts of S0 galaxies, implying that they contain dynamically-cool stellar disks that make up an appreciable fraction of the total mass. If confirmed kinematically, the presence of an outer stellar disk in the majority of these systems would strongly support gas dissipation as the last significant step in their formation history. For individual objects, detailed photometric and kinematic studies have shown indeed that dynamically fragile stellar disks are present in several elliptical galaxies (*e.g.* Rix & White 1992; Scorza & Bender 1995).

In this paper we attempt to move beyond questions of individual misclassification of galaxies, by asking whether there is any significant fraction of “faint elliptical galaxies” that have reached their present state through largely dissipationless merging as the last formation step? To this end we present and discuss kinematic measurements (to $R > R_e$) for a photometrically selected sample of seven bona-fide ellipticals with $M_B \geq -19.5$.

2. Sample and Observations

Our seven galaxies are a random sub-sample of the RC3 catalog (de Vaucouleurs *et al.* 1991), with Hubble types $T \leq -4$ and luminosities below $M_B = -19.5$. No prior kinematic information was used, and nothing, save the luminosity cut, should have biased the selection process towards rapidly rotating objects. We added three S0 galaxies ($T = -2, -3$)

for comparison. The spectra were obtained during two different runs (February 11-14, 1997, and September 29 – October 2, 1997) at the KPNO 4-m telescope, using the RC spectrograph with the KPC-24 grating in second order. In February 1997 the detector was T2KB CCD with 2048^2 pixels of $24\mu\text{m}^2$, which were binned along the slit to yield $1.38''/\text{pixel}$. In the September 1997 run we used a $3\text{k}\times 1\text{k}$ F3KB CCD with $15\mu\text{m}^2$ pixels binned to $0.86''/\text{pixel}$ along the slit. With a $2.5''$ wide slit and on-chip binning by two in the spectral direction the effective instrumental resolution was $\sigma_{instr} \approx 50 \text{ km/s}$. The spectra were centered on $5150\text{\AA}$, covering $\text{H}\beta$ $4861\text{\AA}$, $[\text{OIII}]$ $5007\text{\AA}$, Mg_b $5175\text{\AA}$, Fe $5270\text{\AA}$ and Fe 5335 . Table 1 lists relevant properties of the sample members and the observations, including the adopted major axis position angle and the exposure times. Spectra of several K giants were acquired with the same instrumental setup, and used as kinematic templates. The basic data reduction (bias and dark subtraction, flat-fielding, correction for slit-vignetting, wavelength calibration, airmass and Galactic extinction corrections, correction for instrumental response) was performed using IRAF and MIDAS software.

The stellar kinematics were obtained with the template fitting method, described in Rix & White (1992), after the data were binned along the slit to constant signal-to-noise. Then the best fit mean velocity V and dispersion σ were determined by χ^2 -minimization, along with the best fitting composite stellar template. The resulting error bars are the formal uncertainties based on the known sources of noise. External tests on this and other data sets demonstrate that they are a good approximation to the true uncertainties.

3. Results and Discussion

Figure 1 shows the major and minor axis kinematics for the sample galaxies. Similar to S0 galaxies, the velocity dispersions in these ellipticals are found to drop from a central peak to near our instrumental resolution at large radii. Most sample members exhibit little - if any - rotation along the minor axis, in contrast to their strong rotation along the major axis. Their kinematics suggest that these systems are nearly axisymmetric and likely have a rather simple dynamical structure. Only NGC1588 shows considerable minor axis rotation over the same radial region where the photometry shows isophotal twisting; this is possibly due to the interaction with its close companion (NGC1589).

In Figure 2 we quantify the degree of rotational support in these galaxies, by plotting V/σ as a function of the galactocentric distance along the major axis. As a diagnostic of the dynamical state beyond the effective radius, we adopt the maximum, $(V/\sigma)_{max}$, and the outermost value $(V/\sigma)_{out}$, of each V/σ curve [see Table 1]. In terms of these quantities, Figure 3 re-states the main result: for four out of seven ellipticals, $(V/\sigma)_{max} \gtrsim 2$; two

additional ellipticals have $(V/\sigma)_{max} \gtrsim 1.5$.

As a benchmark, we compare the observed $(V/\sigma)_{max}$ [and $(V/\sigma)_{out}$] values to maximally rotating, oblate “Jeans models.” We consider models where the residual velocities are isotropic in the co-rotating frame. The models are built following the method outlined in Binney, Davies, & Illingworth (1990; see also Carollo & Danziger 1994 for particulars), assuming a Jaffe (1983) law. The curves in Figure 3 represent maximum V/σ within $2R_e$, as a function of projected axis ratio (or inclination). We considered *intrinsic* axis-ratios of $c/a = 0.6$ and $c/a = 0.4$ and models with and without dark halos (which are three times more extended and more massive than the luminous component). As Figure 3 illustrates, no model can reproduce values of $(V/\sigma)_{max} \sim 2$ for any viewing angle, unless it has $c/a \lesssim 0.4$! This implies that all sample galaxies are intrinsically *very* flat. A possible exception is the one distorted galaxy, NGC1588, for which our kinematics are compatible with $c/a < 0.6$, if $(V/\sigma)_{out}$ is considered instead of $(V/\sigma)_{max}$. The probability that 6 out of 7 ellipticals are that flat is vanishing ($\approx 2.4 \times 10^{-6}$) on the basis of the expected shape distribution of low-luminosity oblate spheroidals (Tremblay & Merritt 1996).

Dissipationless simulations of equal mass galaxy mergers mostly produce remnants with $(V/\sigma)_{max} \lesssim 1$ inside $\sim 2R_e$ (*e.g.* Heyl, Hernquist and Spergel, 1996). In addition, for most spin-orbit geometries major mergers also produce significant kinematic misalignment, which would be reflected in minor axis rotation. Neither property is consistent with our sample galaxies, ruling out formation through mergers of nearly equal mass. Unequal mass mergers ($\sim 3 : 1$) can produce remnants with rather disk-like shapes and kinematics (Barnes, 1996; Bekki 1998; Barnes 1998). Barnes (1998) finds remnants to be close to axisymmetric with little minor axis rotation, in this respect consistent with our sample properties. He quantifies the rotational support of the merger remnants by the parameter λ' , the total angular momentum of the most tightly bound half of the stars, normalized by the value for perfect spin alignment. For Barnes’ eight 3:1 mergers $\lambda'_{sim} = 0.38$ with a scatter of 0.04. As λ'_{sim} is not observable, we attempt to construct an analogous quantity $\langle \lambda' \rangle_{obs}$ for our sample members (see Table 1), by estimating the azimuthal velocity, normalized by the local circular velocity and averaged over the inner 50% of the light. For this estimate we had to make the following assumptions: (i) the overall potential is logarithmic; (ii) the velocity dispersions are isotropic; (iii) the galaxy is axisymmetric with an intrinsic axis ratio of 0.4 (or 0.6; see above); (iv) the circular velocity is estimated as $v_c \approx \sqrt{v_\phi^2 + 2\sigma^2}$, which is both correct for non-rotating systems in logarithmic potentials and in the “asymmetric drift” limit for $\rho_* \propto r^{-2}$; (v) the stars are on average 30° from the mid-plane. With this, we find $\langle \lambda' \rangle_{obs} = 0.55$ with a scatter of 0.06 (see Table 1): every observed sample galaxy has considerably more ordered motions than any of the 3:1 mergers. We checked that plausible changes in the above model assumptions will alter not $\langle \lambda' \rangle_{obs}$ by $\langle \lambda' \rangle_{obs} - \lambda'_{sim}$.

The degree of streaming motion found in our sample galaxies is nearly that of the dominant disks of S0 galaxies [*e.g.* Fisher 1997 and the $(V/\sigma)_{max}$ values for the S0s of our sample]. Without exception the sample members appear even more rotationally supported than simulated unequal mass (3:1) mergers (Barnes 1998). This strongly suggests that the dissipative formation of a massive and extended stellar disk has been the last major step in building these galaxies. This inference does not preclude, however, that for some fraction this disk has been heated considerably by subsequent gravitational interactions.

Our data, therefore allow us to push the long-known result that *rotation is important in low-luminosity ellipticals* (Davies *et al.* 1982) one step further: most of these “elliptical” galaxies in an effectively luminosity-limited sample, actually contain stellar disks at large radii, comparable in mass and size to S0s. These kinematic results, combined with earlier photometric evidence (*e.g.*, Rix & White 1990 and references therein), now put this idea on solid observational grounds. Perhaps these results imply that at smaller mass scales the epoch of mergers ended before the epoch of star-formation.

HWR is supported by the Alfred P. Sloan Foundation. CMC is supported by NASA through the grant HF-1079.01-96a awarded by the Space Telescope Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA under contract NAS 5-26555.

REFERENCES

- Bender, R., Surma, P., Doebereiner, S., Moellenhoff, C., Madejsky, R., 1989, A&A, 217, 35
- Barnes, J. 1996 in “Formation of the Galactic Halo ... Inside Out”, eds. H. Morrison and A. Sarajedini, ASP Conference Series, p. 415.
- Barnes, J. 1998 in “Galaxies: Interactions and Induced Star Formation”, Kennicutt, R., Schweizer, F and Barnes, J., Saas Fe Advanced Course 26, Springer, §7.
- Bekki, K. 1998, ApJ, 502, L133.
- Binney, J.J., Davies, R.L., Illingworth, G.D., 1990, ApJ, 361, 78
- Carollo, C.M., Danziger, I.J., 1994, MNRAS, 270, 743
- de Zeeuw, P. T. and Franx, M., 1991, ARA&A, 29, 239
- Davies, R.L., Efstathiou, G., Fall, S.M., Illingworth, G., Schechter, P., 1983, ApJ, 266, 41
- de Vaucouleurs, G., de Vaucouleurs A., Corwin H. G. Jr., Buta R. J., Paturel G., Fouqué P., 1991, Third Reference Catalog of Bright Galaxies, Springer (RC3)
- Fisher, D., 1997, AJ, 113, 950
- Heyl, J.S., Hernquist, L., Spergel, D.N., 1996, ApJ, 463, 69
- Jaffe, W., 1983, MNRAS, 202, 995
- Kormendy, J. and Djorgovski, G., 1989, ARA&A, 27, 235.
- Rix, H-W., White, S.D.M., 1990, ApJ, 362, 52
- Rix, H-W., White, S.D.M., 1992, MNRAS, 254, 389
- Scorza, C., Bender, R., 1995, A&A, 293, 20
- Tremblay, B., Merritt, D., 1996, AJ, 111, 2243
- Weil, M.L., Hernquist, L., 1996, ApJ, 460, 101








Fig. 1.— Rotation velocity V and velocity dispersion σ (both in km/s) along the principal axes as a function of radius (in seconds of arc) for the sample galaxies. Vertical dotted lines are plotted at the half-light radii.






Fig. 2.— The ratio of rotation velocity and velocity dispersion, V/σ , as a function of radius (in arcseconds), illustrating the strong outward increase in rotation support for almost all sample members.




Fig. 3.— Comparison of $(V/\sigma)_{\max}$ (filled squares for the ellipticals; stars for the lenticular galaxies) with oblate, $f(E, J_z)$ Jeans models (thick lines for an intrinsic $c/a = 0.6$, and thin lines for an intrinsic $c/a = 0.4$). The dotted lines indicate models with no dark matter; dashed lines represent models with round dark halos; the solid lines indicate dark halos as flattened as the luminous component. All halos are three times more extended and more massive than the luminous component. Arrows point to the corresponding values of $(V/\sigma)_{\text{out}}$, when these differ from $(V/\sigma)_{\max}$. The typical errorbar on $(V/\sigma)_{\max}$ [and $(V/\sigma)_{\text{out}}$] is given in the upper-right corner.

Name	T	V_{hel} (km/s)	D (Mpc)	M_B (mag)	R_e (arcs)	ϵ	Run	$T_{maj/min}$ (secs)	PA (deg)	$(V/\sigma)_{out}$	$(V/\sigma)_{max}$	$\langle\lambda'\rangle$
NGC386	-5	5707	76	-19.2	11.8 $\ddagger$	0.13	Sep97	18000/8100	5	1.0	1.0	0.47 (0.43)
NGC1588	-4.6	3397	19 $\dagger$	-17.7	12.8	0.30	Sep97	10800/3400	35	1.05	1.4	0.58(0.54)
NGC1603	-5?	5038	67	-19.5	12.2	0.30	Sep97	18000/2100	50	1.5	1.5	0.50 (0.46)
NGC2592	-5	1925	26	-18.9	10.5 $\ddagger$	0.17	Feb97	5400/5400	53	3.0	3.0	0.65 (0.62)
NGC2699	-5	1660	22	-18.1	9.0 $\ddagger$	0.07	Feb97	5400/5400	45	2.1	2.1	0.63 (0.60)
NGC2778	-5	1991	27	-18.8	15.7	0.28	Feb97	5400/3600	40	2.1	2.4	0.54 (0.50)
NGC3605	-5	581	14 $\dagger$	-17.6	21.2	0.33	Feb97	4400/5400	22	1.8	1.8	0.47 (0.43)
NGC2577	-3	2068	28	-19.0	24.3 $\ddagger$	0.38	Feb97	5400/3600	105	3.0	3.8	0.71 (0.68)
NGC3156	-2	980	15 $\dagger$	-17.9	14.4	0.42	Feb97	5400/5400	47	2.4	3.0	0.60 (0.57)
NGC7617	-2	4176	56	-19.1	10.6	0.35	Sep97	18000/12560	30	1.5	1.5	0.62 (0.58)

Table 1: Parameters and observing log for the seven elliptical and three S0 galaxies of our sample. Morphological types T , magnitudes and (outer) ellipticities ϵ are from the RC3 or UGC Catalogs. The half-light radius R_e is from the RC3 catalog, except for the galaxies identified with the symbol $\ddagger$, for which this measurement is derived from our spectral data (since it is not provided by RC3). Absolute magnitudes are corrected for Galactic extinction. We assume $H_o = 75$ km/s/Mpc. Distances are computed directly from the Hubble flow velocities (RC3), except for those galaxies identified by the $\dagger$. For these, the peculiar motion corrected distances from Faber et al. (1989) or Bender et al. (1992) are adopted. $T_{maj/min}$ gives the major and minor axis spectra total exposure times, respectively. PA gives the adopted values for the position angle of the major axis. The columns $(V/\sigma)_{out}$ and $(V/\sigma)_{max}$ list respectively the outermost and the maximum value of the V/σ ratio over the radial range sampled by the data. The spin parameter $\langle\lambda\rangle$ (for $q_{intr} = 0.4$) is defined in the text; values in parentheses refer to $q_{intr} = 0.6$.